


# CPR and First Aid



## ACROSS

- 1 The carotid pulse is checked for \_\_\_\_\_ seconds to determine if the patient's heart is beating.
- 7 In 2-rescuer CPR on an infant, \_\_\_\_\_ compressions are given followed by 2 breaths.
- 9 In child CPR, two \_\_\_\_\_ breaths are given, rather than 2 full breaths (as in adult CPR).
- 10 If the patient has no pulse, you would give \_\_\_\_\_ chest compressions.
- 11 The \_\_\_\_\_ step in adult CPR is to lightly shake the pt. and ask if he/she is OK

## DOWN

- 1 Infant CPR is performed on an infant/child weighing less than \_\_\_\_\_ pounds.
- 2 The head tilt/chin lift is performed, and then you would check for \_\_\_\_\_.
- 3 The compression rate in adult CPR is a \_\_\_\_\_ compressions per minute.
- 4 The second step in adult CPR is to \_\_\_\_\_ 911.
- 5 Adult CPR is performed on a child eight years or older, or on a child weighing more than \_\_\_\_\_ pounds.
- 6 In infant CPR, check the \_\_\_\_\_ pulse for 15 seconds.
- 8 If the patient is not breathing, give \_\_\_\_\_ full breaths before checking his/her pulse.

# CPR and First Aid

