


# Psychiatry 2



## ACROSS

- 1 The initials for the safest type of antidepressant medication is \_\_\_\_\_.
- 7 The first Kubler-Ross Stage of Dying should be \_\_\_\_\_.
- 8 \_\_\_\_\_ grief lasts greater than 1 year, and has symptoms of delusions and hallucinations.
- 11 This type of major depression is found in the elderly, and often occurs in Winter.
- 12 \_\_\_\_\_ are fixed, false beliefs, lasting at least 1 month.
- 13 A teenager who throws a temper-tantrum when she doesn't get her way is \_\_\_\_\_-out.

## DOWN

- 1 Symptoms of this disease include auditory or visual hallucinations, delusions, echolalia, and blunting.
- 2 Avoiding awareness of some painful aspect of reality is called \_\_\_\_\_.
- 3 A severely obese person who can't stop eating, ultimately becomes a chef. This is an example of \_\_\_\_\_ formation.
- 4 \_\_\_\_\_ I Disorder is diagnosed after 1 manic episode that lasts at least 1 week, or requires hospitalization.
- 5 A patient who acts, dresses, and talks like Elvis all day, every day, suffers from \_\_\_\_\_.
- 6 A patient suffering with \_\_\_\_\_ delusions believes that someone else is in love with them.
- 9 Patients who believe that people are all good or all bad suffer from \_\_\_\_\_.
- 10 This type of schizophreniac is immobile and rigid.

# Psychiatry 2

